



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Bill of health was issued to the following-named vessel:

Date.	Vessel.	Destination.	Number of crew.	Number of passengers from this port.	Number of passengers in transit.	Pieces of baggage disinfected.
Sept. 27	Katie.....	Mobile via Belize.	24	.....	.....	.....

*Circular relative to measures against the spread of yellow fever at Zacapa and Gualan.*

(ISSUED BY GOVERNMENT OF GUATEMALA.)

Mosquitoes breed in confined water in cisterns, pipes, etc. These deposits of water should be emptied or covered with a thin layer of petroleum. The water may be used to wash clothes.

In the rooms burn sulphur. This should be done in the morning. At night burn insect powder before retiring.

Use mosquito bars.

Persons who are not immune, recently arrived, etc., should avoid houses where the fever has already occurred and seek houses where it has not been present.

All fever cases are infectious during the first three days. The patients should be isolated, or at least provided with mosquito bars.

Any person falling ill with a high fever that begins with a chill will do well not to accept the case as one of cold or malaria, but consider himself attacked with yellow fever.

As soon as illness begins 1 grain of calomel should be taken, and afterwards, every two hours, 1 teaspoonful of sulphate of magnesia and cream of tartar, use of this mixture to be continued during the first three days; also one-half gram of salol can be taken every three hours during these three days.

Absolutely nothing should be eaten; only lemonade, tea, and gruel should be taken. Milk and alcoholic drinks are injurious.

In case of pains in the stomach, which nearly always indicate intestinal hemorrhage, 20 drops of laudanum should be taken at one time.

From the time that the first case of yellow fever appears in a non-infected town fifteen days elapse before there are secondary cases. This time should be utilized to kill the mosquitos in the house where the sickness occurred.

HAWAII.

*Plague at Waipahu and Honolulu.*

HONOLULU, October 11, 1905.

WYMAN, Washington:

Two plague, Waipahu, October 9, fatal; 1, Honolulu, October 10.

HOB DY.